


## Gonorrhea Infections in North Carolina Reported Gonorrhea Case Data, 2014



# Reported gonorrhea infections have remained stable over the past few years





#### In 2014

- 14,952 gonorrhea infections were reported in North Carolina.
- North Carolina gonorrhea rates are higher than the national rates, similar to many states in the Southeast (143 vs. 106 per 100,000 population; CDC 2013).


## Disparity by race/ethnicity is particularly large for gonorrhea

## What YOU can do

Provide cultural competency training for your staff and colleagues.

Resources in side bar.


## Gonorrhea Infections by Race/Ethnicity, 2014



Co-infection with HIV (diagnosed prior to or within 30 days of gonorrhea infection) has doubled among men in the past few years

## Gonorrhea and HIV Co-infection by Gender, 2005-2014

- The percentage of men who are co-infected with gonorrhea and HIV has been increasing since 2007.
- The percentage of co-infected women remains stable and fairly low.



# Want More Information?

HIV/STD Facts and Figures website: http://

epi.publichealth.nc.go v/cd/stds/figures.html

Centers for Disease Control and Prevention (CDC) Fact Sheet on Gonorrhea:

http://www.cdc.gov/ std/gonorrhea/stdfactgonorrhea.htm

Cultural Competency (C3) Trainings and Information on Care and Prevention in the United States (CAPUS) in North Carolina web site:

http:/ www.med.unc.edu/ ncaidstraining/capus

### **Contact Us**

North Carolina DHHS Communicable Disease Branch

**Phone:** (919) 733-3419

Mailing Address: Communicable Disease Branch Epidemiology Section 1902 Mail Service Center Raleigh NC 27699-1902

Created by the HIV/STD Surveillance Unit 11/16/2015



## Gonorrhea Infections in North Carolina Reported Gonorrhea Case Data, 2014



## What is North Carolina doing to decrease gonorrhea infections?


- In September 2014, the SLPH increased the age cut off for routine gonorrhea screening in women from ≤24 years of age to ≤25 years of age.
- North Carolina provides funds for gonorrhea screening for all women who are seen in a
  publically funded health care facility, such as local health departments and family
  planning settings.
- Two counties in North Carolina are participating in a pilot project to offer partner services to men who report sex with men (MSM) and men with HIV who are diagnosed with gonorrhea.
- The Department of Public Health is moving toward providing testing for gonorrhea at extragenital (rectal and oropharyngeal) sites for men reporting sex with men (MSM) and transgendered people.

## What CLINICIANS can do

- Routinely ask patients about their sexual activity and test those that are sexually active
- Provide treatment consistent with the CDC guidelines (link in side bar)
- Refer partners for evaluation and treatment
- Report cases to the local health department

# In 2014, 75% of all gonorrhea infections were treated correctly in North Carolina

## North Carolina Gonorrhea Infection Treatments in 2014



#### **Treatment Options**

## What YOU can do

If you have a gonorrhea infection, ensure that you and your partners get treatment. Untreated gonorrhea can lead to severe health outcomes, including increased risk for HIV and pelvic inflammatory disease (PID).

- treating gonorrhea is very important, particularly because of the rise of drug resistance.
  - In 2016, the North
    Carolina Division of
    Public Health
    Technical Assistance
    and Training
    Program (TATP)
    nurses will support
    county work to
    improve adherence
    to treatment
    guidelines and
    ensure the correct
    treatments are
    being delivered.

## Recommendations from the CDC 2015 STD Treatment Guidelines:

## Dual Therapy for Uncomplicated Gonococcal Infections Recommended Regimens:

- Ceftriaxone (250 mg IM in a single dose PLUS
- Azithromycin (1 g orally in a single dose ).

Alternative Regimens (where ceftriaxone is not available):

- Cefixime (400 mg orally in a single dose PLUS
- Azithromycin (1 g orally in a single dose).
- Gentamicin (240 mg IM one dose)
   PLUS
- Azithromycin (2 g orally in a single dose).

#### **Data Source:**

North Carolina Electronic Disease Surveillance System (NC EDSS) (data as of July 6, 2015).

State of North Carolina •
Pat McCrory, Governor
Department of Health and
Human Services •
Richard O. Brajer, Secretary
Division of Public Health •
Megan Davies, M.D., Acting
State Health Director
HIV/STD Surveillance Unit
www.ncdhhs.gov •
www.publichealth.nc.gov

N.C. DHHS is an equal opportunity employer and provider.

Created by the HIV/STD Surveillance Unit 11/16/2015